

RENAL IMPAIRMENT DOSING TABLE: ANTIFUNGALS V2.0

Renal Impairment Prescribing Table – ANTIFUNGALS				
Dose adjustments recommended in this table are applicable to the infections detailed on www.antibioticprescribing.ie for the treatment of community infections only. All doses are oral and for adults unless otherwise stated. For patients on dialysis, seek advice from renal team.				
Use either eGFR or calculated CrCl figure to direct to relevant dosing column in tables below.				
Drug (oral unless otherwise stated)	Stage 3A	Stage 3B	Stage 4	Stage 5
	eGFR (ml/min/1.73m ²) or calculated CrCl (ml/min)			
	30 - 50		10 - 30	<10
Fluconazole	No dose adjustments necessary for single dose regimens. 50% dose reduction after initial dose for daily or weekly dosing regimens ≥50mg daily.			
Fluconazole ♦ 150mg stat	No adjustment required			
Fluconazole ♦ 150mg day 1, 4 and 7 then weekly for 6 months				
Fluconazole ♦ 300mg weekly for 2 weeks				
Fluconazole ♦ 50mg every 24 hours	50mg every 48 hours			
Griseofulvin	No adjustment required			
Itraconazole	30-60: No adjustment required		<30: Avoid. Seek specialist advice for alternative	
Terbinafine	<50: 250mg on alternate days			