

# HSE Venous Thromboembolism (VTE) Interim Clinical Guidance for Acute Hospitals during the COVID-19 pandemic

**VTE Protocol Development Group (version 1):** C Kirke (Clinical Lead, Medication Safety Programme, National QI Team), F Ni Áinle (Consultant Haematologist, Rotunda and Mater), D O’Keeffe (Consultant Haematologist, University Hospital Limerick), C Bergin (Clinical Lead, Infectious Diseases, HSE and Consultant in Infectious Diseases, St James’s), V Hamilton (National Clinical Advisory and Group Lead, HSE Acute Operations and Consultant Anaesthetist, University Hospital Waterford), M Pate, A Cronin (Medication Safety Programme, HSE National QI Team).

**VTE Protocol Development Group (update, version 2):** C Kirke, M Pate, G Molloy (NMedication Safety Programme, National Quality & Patient Safety Directorate), F Ni Áinle (Consultant Haematologist, Rotunda and Mater), N O’Connell (Consultant Haematologist, St James’s Hospital), D O’Keeffe (Consultant Haematologist, UHL), C Bergin (Clinical Lead, Infectious Diseases, HSE and Consultant in Infectious Diseases, St James’s), M O’Connor (NCAGL, HSE Acute Operations and Consultant Geriatrician, CUH), A Nichol (Consultant in Critical Care Medicine, SVUH), N O’Hanlon (Chief II Pharmacist, Medicines Information, SVUH), B Cleary (Chief Pharmacist, Rotunda Hospital).

**VTE Patient Information Development Group:** AM O’Neill (CEO, Thrombosis Ireland), C Kirke, M Pate, A Cronin, F Ni Áinle.

**Purpose:** This guidance is to clarify current recommendations for prevention of VTE and patient information for hospitalised adult medical or COVID-19 patients.

**Target Audience:** All healthcare professionals in acute hospitals (protocol and patient information). All patients in acute hospitals (patient information).

## Background

Venous thromboembolism (VTE) or blood clots led to or occurred during hospitalisation of 8,589 people in acute public hospitals in 2020. Up to 70% of blood clots associated with hospitalisation are considered to be potentially preventable with appropriate prophylaxis depending on patient and disease factors. COVID-19 is associated with an increased risk of VTE in hospitalised patients.

## Recommendations

The HSE VTE protocol template includes recommendations for adult patients admitted to acute hospitals with COVID-19, together with all adult medical patients. The template has been updated in April 2022 and should replace previous versions in use.

**All adult patients admitted to hospital**, including patients with COVID-19, require:

- Risk assessment of the patient’s risk of VTE and bleeding, with VTE prophylaxis prescribed and administered as appropriate.
- Patient information (including provision of HSE/Thrombosis Ireland VTE alert card).

**For patients deemed to be at high VTE risk** (due to COVID-19 or Padua Prediction Score of 4 or greater), we suggest:

- **Prophylactic-intensity** low molecular weight heparin (LMWH) or unfractionated heparin (unless contraindicated due to bleeding risk), if the patient has:

- Severe COVID-19 infection (requiring critical care level treatment including high flow oxygen, invasive or non-invasive ventilation, vasopressors),
- Mild-moderate COVID-19 infection who do not have oxygen saturations of 93% or less on room air or low-flow oxygen, and who do not have an elevated bleeding risk. This may include patients who test positive for COVID-19, who require admission from other medical illnesses,
- Medical patients with no evidence of COVID-19.

- **Therapeutic-intensity LMWH\*** *may be considered* in patients admitted to hospital because of moderate COVID-19\*\* AND who have a low bleeding risk *\*for the purpose of reducing risk of death, invasive mechanical ventilation, non-invasive mechanical ventilation, or admission to ICU*

\*\*Moderate COVID-19 is defined as follows: admission to hospital ward level of care (ie not to ICU), not already mechanically ventilated, and not imminently requiring mechanical ventilation or critical care. We suggest that this therapeutic strategy be limited to patients who have oxygen saturations of  $\leq 93\%$  on room air due to COVID-19, or who require low-flow oxygen via nasal prongs or face mask to maintain normal oxygen saturations.


This suggestion also applies to patients who are admitted for another reason but who progress to develop moderate COVID-19.

Patients with COVID-19 should be re-assessed regularly, with VTE prevention adjusted if their condition or level of care changes.

Medical patients and patients with COVID-19 assessed to be high-risk for VTE but with contra-indications to LMWH or heparin should receive mechanical VTE prophylaxis unless contra-indicated. Measure and select the appropriate size(s). Assess fit, compliance and skin integrity at least daily.

## Patient information

Hospitals are asked to ensure patients in all high-risk groups receive the information in these cards.



**For more information:**  
[www.thrombosis.ie](http://www.thrombosis.ie)

**BLOOD CLOT ALERT CARD**

**WHAT IS A BLOOD CLOT?**  
This is the formation of a clot inside a blood vessel, usually in the leg, which may break off and go to the lungs. This can be fatal.

**60%** of clots happen in **HOSPITAL** or in the **90 DAYS** following admission.

Blood clots can be very serious - but there are effective treatments to deal with them and help prevent them

**SIGNS AND SYMPTOMS OF A BLOOD CLOT**


- Swelling or pain in one leg or arm
- Warmth or redness in the leg or arm
- Short of breath or rapid breathing
- Chest pain (particularly when breathing deeply)
- Coughing or coughing up blood

**If you have one or more of these, you may have a clot and need urgent treatment**

Thrombosis Ireland  
Spot The Signs... Save A Life

CSN: 20154240

Seirbhís Stáinte Níos Fearr á Forbairt | Building a Better Health Service



**BLOOD CLOT ALERT CARD**

**Am I at risk?**

**WHAT CAN I DO TO HELP MYSELF?**

- Ask for your risk of blood clots to be assessed, especially if you are in one of the higher risk groups listed opposite
- Walk and move as much as possible
- Drink plenty of fluids
- If directed to use stockings or medication to prevent or treat a clot follow instructions exactly
- Remember, a blood clot in the veins is more likely up to 90 days **after** being in hospital
- If you have any signs or symptoms of a clot, take **immediate action** to seek medical help

You have a **HIGHER CHANCE** of getting a clot in **HOSPITAL** than on an **AEROPLANE!**

**YOU MAY BE AT HIGHER RISK IF YOU:**

- are admitted to hospital and for 90 days after you go home
- have active cancer or receiving cancer treatment
- are pregnant or have had a baby less than 6 weeks ago
- become immobile (more than 3 days in bed / travel non-stop more than 6 hours / in a leg cast)

**RISK MAY INCREASE FURTHER IF:**

- you or a close relative had a blood clot
- you had surgery in the last 90 days
- you have thrombophilia (tendency to clot)
- you are on the oral contraceptive pill or HRT
- you have heart, lung or inflammatory disease
- you are over 60 years of age or are overweight
- you have varicose veins that become red and sore

Blood clot alert cards are available to download from [www.safermeds.ie](http://www.safermeds.ie) and supplies of alert cards are available from [safermeds@hse.ie](mailto:safermeds@hse.ie). Patient and public information as well as alert cards are available translated into multiple languages from [www.thrombosisireland.ie](http://www.thrombosisireland.ie).

# VTE Prevention Protocol

## for In-Patients aged 16 or Over with COVID-19 or Medical Conditions

All hospitalized patients are at increased risk for VTE. VTE is associated with increased morbidity and mortality. Appropriate VTE prevention reduces risk for patients at high risk of VTE.

**Assess all patients as soon as possible (within 14 hours) after the decision to admit. Reassess at consultant review and if clinical condition changes.**

Step 1: VTE risk assessment VTE risk factors	Padua score	VTE risk factors continued	Padua score
Confirmed or presumed COVID-19		At risk, proceed to step 2	
Medical in-patient without a COVID-19 diagnosis		Assess according to Padua Prediction Score (below)	
Immobility expected for at least 3 days (confined to bed +/- bathroom)	3	Active cancer or treatment (chemo- or radiotherapy within 6 months or metastases)	3
Previous DVT/PE	3	Thrombophilia	3
Trauma and/or surgery in previous 30 days	2	Ischaemic stroke (discuss with stroke team) or Acute MI	1
Heart and/or respiratory failure	1	Aged 70 or over	1
Taking oestrogen-containing contraceptive or oral HRT	1	Acute infection or Acute or chronic rheumatologic disorder	1
BMI 30 or greater (obese)	1	Pregnant or up to 6 weeks post-partum*	4*

**Patients with COVID-19: all patients are at risk of VTE; proceed to step 2.**

**Medical patients: Padua Prediction Score 4 or greater = at risk of VTE; proceed to step 2.**

**Padua Prediction Score 3 or less = at low risk of VTE; no prophylaxis required.**

**\*Pregnant or post-partum: Medical admission = at risk of VTE, proceed to step 2. Maternity = follow maternity guidance.**

Step 2: Bleeding risk assessment. Any risk factor below = contra-indication to low molecular weight heparin (LMWH) or heparin	
Active bleeding	On anticoagulant at therapeutic levels/dose, e.g. warfarin, dabigatran, rivaroxaban, edoxaban, apixaban, heparin, enoxaparin: No additional prophylaxis except while anticoagulant held
Platelets less than $50 \times 10^9/L$	Undergoing procedure with high bleeding risk, e.g. neurosurgery, spinal or eye surgery
Bleeding disorder, e.g. haemophilia, Von Willebrand's	History of Heparin-Induced Thrombocytopenia (HIT): Contact haematology or pharmacy
Acquired bleeding disorder e.g. liver failure with PT over 15	Other bleeding risk: if risk of VTE outweighs bleeding risk, consider pharmacological prophylaxis.
Acute stroke (discuss with stroke team)	If risk of bleeding outweighs risk of VTE, consider mechanical VTE prophylaxis
Blood pressure 230 systolic or 120 diastolic or greater	Note: Dual antiplatelet therapy does not preclude prophylactic dose LMWH. There is a lack of data to support therapeutic dose LMWH in patients with COVID-19 who are receiving dual antiplatelet therapy. Consider prophylactic-intensity anticoagulation as an alternative
Epidural or spinal or lumbar puncture in last 4 hours or expected in next 12 hours	

### Step 3: Recommended VTE prevention (or anticoagulation as a "therapeutic" strategy for COVID-19). These recommendations should not be used to guide care for patients with acute venous thromboembolism.

All patients	Adequate hydration, early mobilisation			
Pharmacological	Weight 50-100 kg and GFR over 30 mL/min	Weight 101-150 kg	Weight less than 50 kg	GFR less than 30 mL/min
Prophylactic intensity anticoagulation: All COVID-19 patients not on therapeutic anticoagulation or Medical patients with Padua score $\geq 4$ <b>And</b> No C/I to heparin	Tinzaparin 4500 units once daily or Enoxaparin 40 mg once daily	Tinzaparin 4500 units bd or Enoxaparin 40 mg bd	Tinzaparin 3500 units once daily or Enoxaparin 20mg once daily	Heparin 5000 units twice daily or Tinzaparin 3500 units daily (caution) or enoxaparin 20 mg daily (contra-indicated in GFR less than 15 mL/min)
Therapeutic-intensity LMWH* <b>may be considered</b> in patients admitted to hospital because of moderate COVID-19** AND <b>who have a low bleeding risk</b>	Tinzaparin 175 units / kg once daily or Enoxaparin 1 mg/kg twice daily or 1.5 mg/kg once daily			Data supporting the use of therapeutic-intensity anticoagulation for patients with COVID-19 and renal impairment are lacking. We suggest against therapeutic anticoagulation (in the absence of a VTE event), if GFR is less than 30ml/min. Consider prophylactic-intensity LMWH instead.
This also applies to patients admitted for another reason but who progress to develop moderate COVID-19	** Moderate COVID-19 is defined as follows: admission to hospital ward level of care (ie, not to ICU), not already mechanically ventilated, and not imminently requiring mechanical ventilation or critical care. <b>We suggest that this therapeutic strategy be limited to patients who have oxygen saturations of <math>\leq 93\%</math> on room air due to COVID-19, or who require low-flow oxygen via nasal prongs or face mask to maintain normal oxygen saturations.</b>			

#### Mechanical

COVID-19 patient or High-risk medical patient (score 4 or greater) with contra-indication to heparins	Mechanical compression: Anti-embolism stockings* +/- intermittent pneumatic compression devices (IPCD)/ foot pumps * Do not use in suspected or proven peripheral arterial disease, severe dermatitis, massive leg oedema, leg deformity preventing correct fit, peripheral neuropathy, recent skin graft, allergy to fabric or acute stroke. Use caution and clinical judgement if applying stockings over venous ulcers or wounds. Use IPCD if available, particularly in COVID-19 or acute stroke.
Low-risk medical (score 3 or lower)	No heparin or low molecular weight heparin : No mechanical compression

**Duration: local decision; e.g. until low-risk for VTE or until discharged. May consider prolonged prophylaxis on a case-by-case basis.**

**Step 4: Inform the patient about the signs and symptoms of VTE. Prescribe appropriate prophylaxis.**

**Step 5: As part of the discharge plan, give patients (and family members/carers if appropriate) verbal information and the VTE patient alert card. Give those discharged with prophylaxis information about its importance and how to use it effectively and safely and notify their GP.**

## Key references:

1. Barbar, S. et al. *The Padua Prediction Score*. J Thromb Haemost 2010; 8:2450-7
2. UKCPA HAT Committee QA326.2. *Doses of thromboprophylaxis in extremes of body weight*. Feb 2017
3. Summaries of Product Characteristics, [www.hpra.ie](http://www.hpra.ie)
4. The ATTACC, ACTIV-4a, and REMAP-CAP Investigators. *Therapeutic Anticoagulation with Heparin in Critically Ill Patients with Covid-19*. N Engl J Med, 2021. **385**(9): p. 777-789.
5. The ATTACC, ACTIV-4a, and REMAP-CAP Investigators. *Therapeutic Anticoagulation with Heparin in Noncritically Ill Patients with Covid-19*. N Engl J Med, 2021. **385**(9): p. 790-802.
6. Sholzberg, M., et al. *Effectiveness of therapeutic heparin versus prophylactic heparin on death, mechanical ventilation, or intensive care unit admission in moderately ill patients with covid-19 admitted to hospital: RAPID randomised clinical trial*. BMJ, 2021. **375**: p. n2400.
7. Spyropoulos, A.C., et al. *Efficacy and Safety of Therapeutic-Dose Heparin vs Standard Prophylactic or Intermediate-Dose Heparins for Thromboprophylaxis in High-risk Hospitalized Patients With COVID-19: The HEP-COVID Randomized Clinical Trial*. JAMA Intern Med, 2021.
8. Lopes, R.D., et al., *Therapeutic versus prophylactic anticoagulation for patients admitted to hospital with COVID-19 and elevated D-dimer concentration (ACTION): an open-label, multicentre, randomised, controlled trial*. Lancet, 2021. **397**(10291): p. 2253-2263.
9. Sholzberg, M., et al., *Randomized trials of therapeutic heparin for COVID-19: A meta-analysis*. Res Pract Thromb Haemost, 2021. **5**(8): p. e12638.
10. British Thoracic Society. *Guidance on venous thromboembolic disease in patients with Covid-19*. 2021. Available from: <https://www.brit-thoracic.org.uk/document-library/quality-improvement/covid-19/bts-guidance-on-venous-thromboembolic-disease-in-patients-with-covid-19/>
11. National Institute of Health. *Statement on anticoagulation in hospitalized patients*. 2022. Available from <https://www.covid19treatmentguidelines.nih.gov/therapies/statement-on-anticoagulation-in-hospitalized-patients/>
12. Stevens SM, Woller SC, Baumann Kreuziger L, et al. Executive summary: antithrombotic therapy for VTE disease: second update of the CHEST Guideline and Expert Panel Report. Chest. 2021;160(6):2247-2259. doi:10.1016/j.chest.2021.07.056
13. American Society of Hematology. *COVID-19 and VTE/Anticoagulation: Frequently Asked Questions*. 2022. Available from <https://www.hematology.org/covid-19/covid-19-and-vte-anticoagulation>
14. National Institute of Health. *Antithrombotic Therapy in Patients With COVID-19*. 2022. Available from <https://www.covid19treatmentguidelines.nih.gov/therapies/antithrombotic-therapy/>
15. REMAP-CAP Writing Committee for the REMAP-CAP Investigators. Effect of Antiplatelet Therapy on Survival and Organ Support–Free Days in Critically Ill Patients With COVID-19: A Randomized Clinical Trial. JAMA. Published online March 22, 2022. doi:10.1001/jama.2022.2910 <https://jamanetwork.com/journals/jama/fullarticle/2790488>
16. Moores LK, Tritschler T, Brosnahan S, Carrier, M, et al. Thromboprophylaxis in Patients With COVID-19. A Brief Update to the CHEST Guideline and Expert Panel Report. Chest. Published online Feb 12, 2022. DOI: <https://doi.org/10.1016/j.chest.2022.02.006>

Next review: September 2022

Content will be kept under regular review. Please check website for COVID-19 HSE Clinical Guidance and Evidence for most recent update